

Collisionless Relaxation of Stellar Systems

Henry E. Kandrup

Department of Astronomy and Department of Physics and Institute for Fundamental Theory, University of Florida, Gainesville, FL 32611

Abstract. The objective of the work summarised here has been to exploit and extend ideas from plasma physics and accelerator dynamics to formulate a unified description of collisionless relaxation that views violent relaxation, Landau damping, and phase mixing as (manifestations of) a single phenomenon. This approach embraces the fact that the collisionless Boltzmann equation (*CBE*), the basic object of the theory, is an infinite-dimensional Hamiltonian system, with the distribution function f playing the role of the fundamental dynamical variable, and that, interpreted appropriately, an evolution described by the *CBE* is no different fundamentally from an evolution described by any other Hamiltonian system. Equilibrium solutions f_0 correspond to extremal points of the Hamiltonian subject to the constraints associated with Liouville's Theorem. Stable equilibria correspond to energy minima. The evolution of a system out of equilibrium involves (in general nonlinear) phase space oscillations which may – or may not – interfere destructively so as to damp away.

1. Introduction

A satisfactory theory of collisionless relaxation must address two general issues, namely (1) the form of the equilibrium towards which any given set of initial conditions should evolve and (2) the overall efficiency of this approach towards equilibrium or, more generally, the physics that drives the evolution. For the case of real galaxies, these questions will be impacted to at least some extent by dissipation associated, e.g., with gas dynamics and/or discreteness effects, i.e., collisionality. Collisionless relaxation is an idealisation in which these effects are completely ignored.

Work in this area dates back at least to the 1960's, when a number of different workers, including Hénon (1961), King (1962), and Lynden-Bell (1967) argued (1) that there should be a coarse-grained approach *towards* (albeit not necessarily *to*) a unique equilibrium, namely that described by so-called Lynden-Bell statistics, and (2) that this approach should proceed on a dynamical time scale, t_D , since it is a collective process, unlike the theory of collisional relaxation which had been developed by Chandrasekhar some twenty years earlier. Later workers subsequently shifted the focus in a number of different ways, notably by asking specifically which initial conditions give rise to which final equilibria (e.g., Ziegler & Wiechen 1989). However, it is probably fair to say that almost all the

work on collisionless relaxation over the past thirty years has been formulated in the context of ideas that evolved in the 1960's.

The objective of the work described here has been to apply to collisionless self-gravitating systems various ideas and techniques from plasma physics and accelerator dynamics which have proven successful in explaining real experiments, the obvious point being that, unlike galactic dynamics, these disciplines allow for the possibility of controlled experiments whereby theoretical predictions can be confirmed and/or falsified. Section 2 discusses some issues related to the interpretation of the *CBE*, the starting point for any theory of collisionless relaxation. Section 3 then identifies the correct mathematical sense in which, as is generally asserted, the *CBE* is a Hamiltonian system. Section 4 exploits this Hamiltonian formulation to present a coherent description of what happens as a generic initial condition evolves into the future. Section 5 concludes with a discussion of the phase mixing exhibited by *CBE* characteristics, i.e., orbits in the self-consistent potential, focusing in particular on the possible role of chaos.

2. The Collisionless Boltzmann Equation

The basic assumption underlying collisionless relaxation is that the system in question can be described by a one-particle distribution function, $f(\mathbf{x}, \mathbf{v}, t)$, defined as a phase space mass density, the evolution of which is governed by the collisionless Boltzmann equation (*CBE*), which implies free streaming in the self-consistent potential associated with f (Hénon 1982).

It is often asserted that, in the $N \rightarrow \infty$ limit, where collisionality becomes irrelevant, the *CBE* and the N -body problem coincide. However, this correspondence is not completely trivial. For the special case of a singular initial condition,

$$f(\mathbf{x}, \mathbf{v}, 0) = \sum_{i=1}^N m_i \delta_D[\mathbf{x} - \mathbf{x}_i(0)] \delta_D[\mathbf{v} - \mathbf{v}_i(0)], \quad (1)$$

in which point masses are located in phase space with arbitrary precision, a solution to the *CBE* is equivalent mathematically to the full N -body problem. However, such singular solutions are not what interest the theorist exploiting the *CBE*. Rather, he or she is interested typically in the evolution of a smooth initial $f(0)$ or the construction of a smooth equilibrium f_0 which may be given as a function of global isolating integrals (although this is not necessary (Kandrup 1998c)). A crucial piece of the physics thus entails passing from singular to smooth distribution functions, which is decidedly nontrivial. Perhaps the best way to give meaning to such a smooth f is to adopt the tact taken by plasma physicists (e.g., Klimontovich 1980) and cosmologists (Peebles 1980), assuming that any given realisation of the N -body problem, performed either by nature or a computer, entails sampling a smooth $f(0)$ to generate initial conditions for the N -body problem and then evolving these into the future.

However, given this interpretation there is no reason *a priori* to expect smooth *CBE* characteristics, i.e., orbits in the smooth potential associated with f , to have anything to do with N -body orbits. In particular, there is no contradiction between N -body orbits which are chaotic and *CBE* characteristics which are completely integrable. This is well known from ordinary kinetic theory. One

anticipates that orbits in a gas of molecules interacting via short range forces will be chaotic (as has been proved for the case of hard sphere interactions), but the mean field characteristics for a homogeneous system in equilibrium are all integrable constant velocity trajectories.

Nevertheless, much of the galactic dynamicist's intuition is based on orbits in smooth potentials, so that one might hope that there is some average sense in which, at least for finite time intervals, *CBE* characteristics track *N*-body orbits. No hard results about this issue have yet been proved. However, there is at least one conjecture (Kandrup 1998a) which has not yet been disproved: Suppose that one performs an ensemble of different *N*-body simulations, all generated by sampling the same smooth $f(0)$ but with one orbit always starting at the same $(\mathbf{r}(0), \mathbf{v}(0))$. One might then suppose that, at least for large *N*, the *rms* deviation of the *N*-body orbits from the *CBE* characteristic associated with the same initial $(\mathbf{r}(0), \mathbf{v}(0))$ will satisfy

$$\delta r_{rms}(t) \sim F(N)\exp(t/\tau), \quad (2)$$

where $\tau \sim t_D$ is the characteristic Lyapunov time on which solutions to the *N*-body problem diverge, and $F(N)$ is a decreasing function of *N* (see, e.g., Kandrup & Smith 1991, Goodman, Hoggie, & Hut 1994).

3. The Hamiltonian Character of the Evolution

Galactic astronomers are cognizant of the fact that the *CBE* is a Hamiltonian system. However, most probably do not know the exact sense in which this is true. There is the implicit idea that, because *CBE* characteristics correspond to orbits in a (in general) time-dependent Hamiltonian system, the *CBE* must itself be Hamiltonian. However, this is not really the point. Viewing the physics as corresponding to orbits in a time-dependent potential is a cheat since this does not incorporate self-consistency in a fundamental way! Rather, a proper Hamiltonian formulation entails a very different viewpoint in which the distribution function f itself is the fundamental dynamical variable.

Such a Hamiltonian formulation was first presented by Morrison (1980) for the Vlasov equation, the plasma analogue of the *CBE*. Here f is taken as the basic dynamical variable which evolves in the infinite-dimensional phase space of distribution functions. For the case of gravitational interactions, the Hamiltonian

$$\begin{aligned} \mathcal{H}[f] = & \int d^3x d^3v \frac{1}{2} v^2 f(\mathbf{x}, \mathbf{v}, t) \\ & - \frac{G}{2} \int d^3x d^3v \int d^3x' d^3v' \frac{f(\mathbf{x}, \mathbf{v}, t) f(\mathbf{x}', \mathbf{v}', t)}{|\mathbf{x} - \mathbf{x}'|} \end{aligned} \quad (3)$$

is nothing other than the mean field energy, as identified, e.g., by Lynden-Bell & Sanitt 1969). The first term represents the mean kinetic energy and the second represents the gravitational potential energy associated with the mass density $\rho = \int d^3v f$. The *CBE* is most easily written in a Hamiltonian form by generalising the notion of an ordinary Poisson bracket. Specifically, given two functionals of the distribution function f , say $\mathcal{A}[f]$ and $\mathcal{B}[f]$, one can define the

action of a Lie bracket $[\cdot, \cdot]$ as

$$[\mathcal{A}, \mathcal{B}] = \int d^3x d^3v f \left\{ \frac{\delta \mathcal{A}}{\delta f}, \frac{\delta \mathcal{B}}{\delta f} \right\} \quad (4)$$

where $\{ \cdot, \cdot \}$ is the ordinary Poisson bracket and $\delta/\delta f$ is a functional derivative. What is important about this Lie bracket is that, like the Poisson bracket, it is a linear antisymmetric operation that satisfies the Jacobi identity, which implies that it can be used to generate a Hamiltonian evolution (see, e.g., Arnold 1989). The key point then is that, in terms of the Hamiltonian (3), the Lie bracket (4) yields the *CBE* as

$$\frac{\partial f}{\partial t} + [\mathcal{H}, f] = 0. \quad (5)$$

By exploiting the fact that

$$\delta \mathcal{H} / \delta f = E = \frac{1}{2} v^2 + \Phi(\mathbf{x}, t), \quad (6)$$

where E is the energy of a test particle moving in the self-consistent potential $\Phi[f]$, eq. (5) can be rewritten in the standard Poisson bracket form (see, e.g., Binney and Tremaine 1987)

$$\frac{\partial f}{\partial t} - \{E, f\} = 0. \quad (7)$$

One important feature associated with the *CBE* is Liouville's Theorem, which implies the existence of an infinite number of conserved quantities, the so-called Casimirs. Specifically, any function $\chi(f)$ defines a phase space functional

$$C[f] = \int d^3x d^3v \chi(f), \quad (8)$$

the numerical value of which is constant in time, i.e., $dC/dt \equiv 0$. The fact that f satisfies these constraints, which are associated with various internal symmetries (Morrison and Eliezur 1996), implies that its evolution is restricted to a reduced, but still infinite-dimensional, hypersurface in the phase space of distribution functions.

All of this is correct mathematically, but still one might ask: what are the p 's and q 's? What are the canonically conjugate variables corresponding to the coordinates and momenta of ordinary particle mechanics? The short answer to this is that, as for other Hamiltonian theories of continuous media, such as the equations for an incompressible two-dimensional fluid, there is no easy decomposition of the dynamical variables into canonical pairs (see, e.g., Morrison 1998). Once one has restricted attention to a reduced phase space in which the values of all the Casimirs are fixed, Darboux's Theorem (see, e.g., Arnold 1989) implies that, at least locally, such conjugate variables exist. However, identifying them is hard and, moreover, there is no guarantee that they can smoothly cover the entire phase space. This means that, when visualising an evolution governed by the *CBE*, one *can* view the physics as being no different fundamentally from

more familiar Hamiltonian systems, but that one has to visualise the evolution in phase space, rather than configuration space. Given a “feel” for Hamiltonian mechanics as viewed in phase space, and an appreciation of the subtleties that arise when one allows for infinite degrees of freedom (in particular, the notion of phase mixing discussed in the following section), visualising an evolution governed by the *CBE* is really not all that complicated.

To make all this mathematics somewhat less obscure, it is useful to consider a much simpler mechanics problem that incorporates many of the same features, namely the equations of motion for a freely rotating three-dimensional rigid body. Specifically, the Euler equations for solid body rotations constitute (cf. Kandrup 1990) a non-canonical Hamiltonian system with a three-dimensional phase space and an evolution generated by a Lie bracket that is *not* the Poisson bracket of ordinary particle mechanics.

Working in Cartesian coordinates, the three dynamical variables can be taken as the three components of angular momentum, L_i ($i = x, y, z$), which coordinatize a three-dimensional phase space. The Hamiltonian is nothing other than the kinetic energy which, in terms of the principal moments of inertia, takes the form

$$H = \sum_{i=1}^3 \frac{L_i^2}{2I_i}. \quad (9)$$

The Lie bracket corresponds to the generator of the three-dimensional rotation group, for which

$$[A, B] = \sum_i \sum_j \sum_k \epsilon_{ijk} L_k \frac{\partial A}{\partial L_i} \frac{\partial B}{\partial L_j} \equiv \sum_i \sum_j \sum_k J_{ij} \frac{\partial A}{\partial L_i} \frac{\partial B}{\partial L_j}. \quad (10)$$

It is easily verified that this Hamiltonian and this bracket combine to yield equations of motion of the form

$$\dot{L}_i = [L_i, H] = \sum_j \sum_k \epsilon_{ijk} L_k L_j I_j^{-1}, \quad (i = x, y, z) \quad (11)$$

which are nothing other than the ordinary Euler equations.

An evolution governed by these equations is constrained by the existence of a conserved quantity, namely the total squared angular momentum

$$C[L_i] = \sum_{i=1}^3 L_i^2. \quad (12)$$

It follows that only two of the three components of angular momentum can be specified independently, and that the effective phase space is really two-dimensional. In principle, one can extract a canonically conjugate pair of variables, but doing so is not all that easy (or illuminating).

4. The Approach towards Equilibrium

Stated succinctly, an evolution governed by the *CBE* corresponds to oscillations in an infinite-dimensional phase space, oscillations which, in many cases, may

exhibit destructive interference and, consequently, damp away: An infinitesimal perturbation away from a stable equilibrium f_0 will, when evolved into the future, exhibit linear phase space oscillations which may or may not exhibit Landau damping/phase mixing. Nonlinear deviations will exhibit nonlinear oscillations about one or more equilibria. Because the *CBE* is Hamiltonian, an initial $f(0)$ cannot exhibit a pointwise approach towards an equilibrium distribution f_0 . However, one *can* get a phase mixing of (linear or nonlinear) modes which implies that, as probed by the behaviour of such observables as the mass density $\rho(\mathbf{x})$ or the velocity dispersion tensor $\sigma_{ij}(\mathbf{x})$, there is an approach towards a time-independent f_0 (Kandrup 1998b).

The obvious question, of course, is: towards which equilibrium f_0 will some initial $f(0)$ evolve? The key to addressing this issue is the recognition that the *CBE* implies that all equilibrium solutions f_0 are energy extremals. More precisely, one can prove that a distribution function f_0 corresponds to a time-independent solution, for which $\partial f_0 / \partial t \equiv 0$, if and only if the first variation of the Hamiltonian (3) vanishes identically for all perturbations δf that satisfy the constraints associated with Liouville's Theorem, i.e., $\delta^{(1)}\mathcal{H} \equiv 0$. This result, which was first discovered in the context of galactic dynamics by Bartholomew (1971) and subsequently discovered by the plasma physicists nearly two decades later, follows as a direct corollary of the observation that the “dynamically accessible” perturbations which satisfy all the constraints associated with Liouville's Theorem are all related to the original f_0 by a canonical transformation in terms of some generating function h , i.e.,

$$f_0 \rightarrow e^{\{h, \cdot\}} f_0 = f_0 + \{h, f_0\} + \frac{1}{2!} \{h, \{h, f_0\}\} + \dots \quad (13)$$

It is easy to see that, if the energy extremal is a local minimum, so that the second variation $\delta^{(2)}\mathcal{H} \geq 0$ for all linear perturbations, the equilibrium is linearly stable. When subjected to a linear perturbation, the distribution function will execute phase space motions analogous to a particle which, initially at rest at the minimum of some potential, is given an infinitesimal phase space perturbation corresponding to a nonnegative kinetic energy and a nonnegative potential energy relative to the extremal point.

Far less trivial, but also true, is the fact that, for fixed values of all the Casimirs $C[f]$, there is always a global energy minimum. This was first proven directly by Wiechen, Ziegler, & Schindler (1988). Alternatively, the existence of a global minimum follows as a corollary to the proof of global existence first established by Pfaffelmoser (1992). Global existence, i.e., the fact that sufficiently smooth initial data never give rise to such singular behaviour as caustics or shocks, is itself important, and not completely obvious physically. One knows, for example, that global existence does *not* hold for the equations that describe a perfect fluid, even though these equations can be derived from the *CBE* by implementing a truncated moment expansion.

If, for fixed values of the conserved quantities $C[f]$, there exists only one energy extremal, f_0 , that extremal must correspond to a global energy minimum and f_0 must be globally stable. This implies that *any* $f(0)$ generated from f_0 by a canonical transformation which leaves invariant all the constraints associated with Liouville's Theorem will simply evolve so as to execute (in general nonlinear) oscillations about the minimum energy state. Explicit examples of such

globally stable equilibria, which include some of the spherical polytropes, have been constructed by Aly (see, e.g., Aly 1994).

In a more general setting, where the reduced phase space hypersurface admits more than one energy extremal, the physics is more complicated in practice although what is going on is still straightforward in principle. In such settings, the distribution function f will oscillate about one or more energy minima just like a particle in a potential can oscillate about one or more energy minima, each of which corresponds to an extremal point where the kinetic energy vanishes and the potential energy is a local minimum.

But how should one visualise the evolution of a small perturbation away from a collisionless equilibrium? Here the important message is that one ought not to distinguish between Landau damping and phase mixing: Viewed appropriately, Landau damping *is* the phase mixing of a wave packet of normal modes. This was first shown for the case of a homogeneous electrostatic plasma by Case (1959), who first computed a complete set of normal modes for the evolution equation satisfied by a linearised perturbation and then demonstrated that wave packets constructed from these modes damp at the rate that Landau (1946) had originally derived.

It is not, however, true that every linearised perturbation of every collisionless equilibrium will damp! Linear Landau damping is guaranteed if all the normal modes are continuous (i.e., if the linearised evolution equation has no point spectrum), but discrete modes correspond to physical perturbations that need not damp away (Habib, Kandrup, & Yip 1986). As discussed more carefully in Kandrup (1988b), the physics here is analogous to what arises in ordinary quantum mechanics. If one considers an observable like angular momentum with a discrete spectrum, it is possible to construct well behaved eigenstates which, when evolved into the future, maintain their coherence for all times. If, however, one considers observables like position or linear momentum, where the spectrum is continuous, the situation is completely different. In this case, a nonsingular initial state must be constructed from a continuous set of singular eigendistribution, so that the best that one can do is construct a localised wave packet. However, when evolved into the future such a wavepacket will necessarily spread because different eigendistributions have different phase velocities.

Especially given the common intuition that perturbations of a “realistic” plasma will always Landau damp, it is important to stress that there *are* known examples, both in plasma physics and galactic dynamics, of geometries where one can have solutions characterised by undamped oscillations. Perhaps the best known example is provided by the so-called Van Kampen (1955) modes, which arise oftentimes in a plasma when the equilibrium distribution vanishes identically for particle speeds larger than some critical value. First predicted analytically, these modes are well known to experimentalists.

Louis & Gerhart (1988) and Sridhar (1989) have demonstrated explicitly that one can also construct models of spherical galaxies that correspond seemingly to time-independent equilibria perturbed by finite amplitude undamped oscillations. Whether such configurations could arise as a consequence of a realistic, or quasi-realistic, evolution is not clear. However, numerical simulations by Mineau et al (1990) have shown that, at least for the toy model of one-dimensional gravity with infinite plane sheets, it is possible to create such a

pulsating configuration by colliding two reasonable time-independent equilibria. These gravitational examples are more complicated than the simple examples that give rise to Van Kampen modes, but the physics is very similar.

The conventional picture of Landau damping (see, e.g., Stix 1962) interprets the damping as resulting from a resonance between unperturbed particles moving with physical velocity $\mathbf{v}$ and a wave, i.e., the perturbation, propagating with a phase velocity $\mathbf{c}$. The obvious point, then, is that if there are no particles with physical velocity $\mathbf{c}$, a wave travelling with velocity $\mathbf{c}$ has nothing with which to resonate so that it will not damp. Landau’s original derivation of damping via an evaluation of poles in the complex plane is only valid for equilibria that are analytic functions of velocity, which precludes the possibility of velocities for which the unperturbed f_0 vanishes identically.

Modulo nontrivial boundary conditions such as phase space holes, linearised perturbations of a stable equilibrium would be expected to exhibit linear Landau damping; and, by analogy, one might expect that larger amplitude perturbations will exhibit nonlinear Landau damping. However, if one visualises a generic initial $f(0)$ as a large perturbation away from *some* equilibrium, the form of which one need not know, it would appear reasonable to view its possible approach towards equilibrium as a manifestation of nonlinear Landau damping. It thus seems natural to argue that the phenomenon that the astronomer is wont to interpret as violent relaxation is really the gravitational analogue of what the plasma physicist terms nonlinear Landau damping.

One potential advantage of such an interpretation is the fact that the words Landau damping suggest an evolution that exhibits significantly more coherence than is usually assumed in violent relaxation. Lynden-Bell’s (1967) original analysis presupposed that the coarse-grained final equilibrium is “random,” subject only to some set of holonomic constraints (including conservation of number and energy) and a coarse-grained version of Liouville’s Theorem, which ensures that the phase space evolution is incompressible. However, numerical simulations by numerous workers (e.g., Van Albada 1982, Quinn & Zurek 1988, Kandrup et al 1994), even those with almost no softening, demonstrate unambiguously that individual particles (and thus, presumably, phase elements in the *CBE*) exhibit a significant remembrance of things past in terms of their binding energies.

5. Regular and Chaotic Phase Mixing

Section 4 focused on the phase mixing exhibited by $f(t)$ as it evolves in the phase space of distribution functions. Also interesting, however, and more consonant with the usual way in which galactic astronomers view collisionless relaxation, is the question of how initially localised phase space elements disperse as a result of an evolution governed by the *CBE*. Honest computations of this sort have not yet been effected numerically. However, some insights into what one might expect can be derived by considering a much simpler problem, namely the mixing exhibited by a phase element evolved in a fixed potential.

What happens if an initially localised phase element (or, equivalently, a collection of points sampling that phase element) is evolved into the future? How fast, and in what sense, will one observe an approach towards some time-independent, or nearly time-independent equilibrium? Does the form of the

evolution depend sensitively on the details of the potential, or does it depend simply on whether the phase element corresponds to chaotic or regular motion?

This problem can be, and has been, investigated by (1) selecting a two- or three-dimensional potential, (2) choosing a small phase space cell of fixed energy, (3) sampling that cell to get a collection of individual initial condition, (4) evolving these initial conditions into the future, and then (5) analysing the statistical properties of the resulting trajectories as a function of time t (Kandrup & Mahon 1994, Mahon et al 1995, Merritt & Valluri 1996, Kandrup 1997). The statistical analysis involved tracking three different diagnostics:

- The convergence of coarse-grained reduced distributions towards time-independent forms. This involved, e.g., binning the data to extract such reduced distributions as $f(y, v_y, t)$ and then determining whether such an $f(t)$ will converge towards a time-independent invariant $f_{eq}(y, v_y)$. Convergence was defined with respect to the “norm”

$$||f(Z_a, Z_b, t) - f_{eq}(Z_a, Z_b)|| = \left(\sum_a \sum_b |f(Z_a, Z_b, t) - f_{eq}(Z_a, Z_b)|^p \right)^{1/p}, \quad (14)$$

for $Z_a \neq Z_b = x, y, z, v_x, v_y$, or v_z and with $p = 1$ or 2 .

- The evolution of moments such as the velocity dispersion σ_{v_x} which, for an initially localised phase element, start small but eventually asymptote towards a much larger value.
- The evolution of other moments such as the mean velocity $\langle v_x \rangle$ which, in many cases, eventually asymptote towards zero.

Investigations of motions in a large number of different potentials have shown that the detailed choice of potential is relatively unimportant, and that the most important distinction is between what Merritt & Valluri have termed *regular* and *chaotic phase mixing*. For regular phase elements, i.e., elements for which the sampled orbits are all regular, the reduced distributions approach an equilibrium or near-equilibrium as a power law in time, i.e.,

$$||f(t) - f_{eq}|| \sim (t/\tau)^{-p}. \quad (15)$$

The growing moments grow as a power law in time until they saturate at a constant value, whereas the decreasing moments damp as a power law in time. Fully chaotic phase elements, i.e., phase elements in two-dimensional potentials with one positive Lyapunov exponent and those in three-dimensional potentials with two positive Lyapunov exponents, evolve very differently. For such phase elements, the reduced distributions approach a (near-)equilibrium exponentially in time, i.e.,

$$||f(t) - f_{eq}|| \sim \exp(-t/\tau), \quad (16)$$

the growing moments initially grow exponentially, and the decreasing moments decrease exponentially. Partially chaotic three-dimensional phase elements, with only one positive Lyapunov exponent, exhibit exponential evolution in some directions and power law evolution in others.

In every case, the natural time scale τ is of order a characteristic dynamical, or crossing, time, but chaotic and regular phase mixing are very different: Exponential evolution is much more efficient than power law evolution. This simple picture is complicated by various technical points, such as the fact that phase mixing can proceed at significantly different rates in different phase space directions and that chaotic mixing can be impeded for surprisingly long times by such obstructions as cantori or an Arnold web (see, e.g., Lichtenberg & Lieberman 1992). However, these are relatively minor perturbations on the basic conclusion: *In generating a well-shuffled state, chaotic mixing is far more efficient than is regular mixing.*

But what is actually seen if one examines visually the distorted phase element or the locations of the points that were evolved from a sampling of the initial $f(0)$? The full four- or six-dimensional phase element becomes stretched and distorted in a volume-preserving fashion which, when projected onto a two-dimensional plane, gives rise to a complicated pattern of tendrils and whorls. The Hamiltonian character of the flow implies that these whorls can never disappear, but the power associated with these structures cascades down to progressively shorter scales. Also interesting, and potentially significant, is the fact that even very weak non-Hamiltonian perturbations, modeled as friction and noise, can be surprisingly efficient in “fuzzing out” these short range structures, thus facilitating a more complete approach towards a near-equilibrium (see, e.g., Habib, Kandrup, & Mahon 1997).

The form of the small scale structures that evolve and the relative efficacy of regular and chaotic phase mixing are evident from Figures 1 and 2, which exhibit the (x, y) coordinates associated with 50625 orbit samplings of initially localised chaotic and regular phase elements in the so-called dihedral potential (cf. Mahon et al 1995). In each case the initial phase element was generated by uniformly sampling a square of side $\Delta y = \Delta v_y = 0.2$ in the $y - v_y$ plane, setting $x = 0$, and then solving for $v_x = v_x(x, y, v_y, E) > 0$, with $E = 1.0$ the energy.

One final point remains to be stressed: *A priori*, whether or not the *CBE* itself is chaotic need have little to do with whether or not *CBE* characteristics are chaotic! The fact that portions of the flow in a self-consistent evolution exhibit chaotic mixing does not imply that the *CBE* is chaotic. Saying that the *CBE* is chaotic should be a statement about the evolution of the distribution function in phase space, not about orbits in the self-consistent potential Φ associated with f .

Acknowledgments. It is a pleasure to acknowledge useful discussions, and arguments, with Paul Channell, Barbara Eckstein, Salman Habib, Ted Kirkpatrick, Donald Lynden-Bell, Elaine Mahon, Dave Merritt, Phil Morrison, Tom O’Neil, Daniel Pfenniger, Ilya Pogorelov, Haywood Smith, and Christos Siopis.

References

- Aly, J. J. 1994, in *Ergodic Concepts in Stellar Dynamics*, V. G. Gurzadyan and D. Pfenniger, New York: Springer, 226
- Arnold, V. I. 1989, *Mathematical Methods of Classical Mechanics*, Berlin: Springer
- Bartholomew, P. 1971, *MNRAS*, 151, 333

- Binney, J. & Tremaine, S. 1987, *Galactic Dynamics*, Princeton: Princeton University Press
- Case, K. 1959, *Ann. Phys. (NY)* 7, 349
- Goodman, J., Heggie, D., & Hut, P. 1994, *ApJ*, 415, 715
- Habib, S., Kandrup, H. E., & Mahon, M. E. 1997, *ApJ* 480, 155
- Habib, S., Kandrup, H. E., & Yip, P. F. 1986, *ApJ*, 309, 176
- Hénon, M. 1961, *Ann. d'Ap.*, 24, 369
- Hénon, M. 1982, *A&A*, 114, 211
- Klimontovich, Yu. 1980, *Kineticheskaya Teoriya Elektromagnitnykh Protsessov*, Moscow: Nauka
- Kandrup, H. E. 1990, *ApJ*, 351, 104
- Kandrup, H. E. 1998a, *Ann. N. Y. Acad. Sci.* 848, 28
- Kandrup, H. E. 1998b, *ApJ*, 500, 120
- Kandrup, H. E. 1998c, *MNRAS*, in press.
- Kandrup, H. E. 1999d, *MNRAS*, in press.
- Kandrup, H. E. & Mahon, M. E. 1994, *Phys. Rev. E* 49, 3735
- Kandrup, H. E., Mahon, M. E., & Smith, H. 1994, *A&A*, 271, 440
- Kandrup, H. & Smith, H. 1991, *ApJ* 374, 255
- King, I. 1962, *AJ*, 67, 471
- Landau, L. D. 1946, *Soviet Phys. – JETP* 10, 25
- Lichtenberg, A. J & Lieberman, M. A. 1992, *Regular and Chaotic Dynamics*, Heidelberg: Springer
- Louis, P. D. and Gerhard, O. E. 1988, *MNRAS*, 233, 337
- Lynden-Bell, D. 1967, *MNRAS*, 136, 101
- Lynden-Bell, D. & Sanitt, N. 1969, *MNRAS*, 143, 167
- Mahon, M. E., Abernathy, R. A., Bradley, B. O., & Kandrup, H. E. 1995, *MNRAS*, 275, 443
- Merritt, D. & Valluri, M. 1996, *ApJ*, 471, 82
- Mineau, P., Feix, M. R. & Rouet, J. L. 1990, *A&A*, 228, 344
- Morrison, P. J. 1980, *Phys. Lett. A* 80, 383
- Morrison, P. J. 1998, *Rev. Mod. Phys.* 70, 467
- Morrison, P. J. & Eliezur, S. 1986, *Phys. Rev. A* 33, 4205
- Peebles, P. J. E. 1980, *The Large Scale Structure of the Universe*, Princeton: Princeton University
- Pfaffelmoser, K. 1992, *J. Diff. Eqn.*, 95, 281
- Quinn, P. J. & Zurek, W. H. 1988, *ApJ* 331, 1
- Sridhar, S. 1989, *MNRAS*, 238, 1159
- Stix, T. H. 1962, *The Theory of Plasma Waves*, New York: McGraw-Hill
- Van Albada, T. S. 1982, *MNRAS*, 201, 939
- Van Kampen, N. G. 1955, *Physica*, 21, 949
- Wiechen H., Ziegler, H., & Schindler, K. 1988, *MNRAS*, 232, 623
- Ziegler, H. J., & Wiechen, H. 1989, *MNRAS*, 238, 1261




Figure 1. The evolution of an initially localised ensemble of 50,625 chaotic orbits in the dihedral potential recorded at times $t = 2, 4, 8$, and 16.




Figure 2. The evolution of an initially localised ensemble of 50,625 regular orbits in the dihedral potential recorded at times $t = 2, 4, 8$, and 16.